

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.


Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS


**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**


Cetylpyridinium Chloride: Butanol  
2:1 by weight




**Figure 1.** Phase diagrams with the surfactant CPC and cosurfactant butanol in the ratio of 2 to 1 by weight. The region enclosed in black line is the one with double the standard molar ratio of structure directing agent (1.2 : 1.0 triethylamine : phosphoric acid). The dash-line region represents standard molar ratio of triethylamine in the aqueous phase (0.6 : 1.0 triethylamine : phosphoric acid). A-D are compositions chosen for hydrothermal synthesis with double the molar ratio of triethylamine. A'-C' are compositions chosen for hydrothermal synthesis with the standard molar ratio of triethylamine.




**Figure 2.** Products from conventional heating (6 hrs at 180°C) with double the standard amount of triethylamine. Figures 2A to 2D correspond to microemulsion compositions A to D respectively from Figure 1. Figure 2E is the control experiment without the microemulsion. The scale bars are 20 microns for A, B, and D; 200 microns for C and E.




**Figure 3.** (A) ~ (E) X-ray diffraction patterns for products shown in Figure 2A to 2E respectively. .




**Figure 4.** Products from microwave heating (17 min at 180°C) with double the standard amount of triethylamine. Figures 4A to 4D correspond to microemulsion compositions A to D respectively from Figure 1. Figure 4E is the control experiment without the microemulsion. The scale bars are A-10µm, B-2µm, C-20µm, D-2 µm, and E-20µm.



**Figure 5.** Products from conventional heating (6 hrs at 180°C) with standard triethylamine concentration. 5A is from a microemulsion with composition C' on Figure 1. 5B is the control experiment without the microemulsion. The scale bars are 200  $\mu\text{m}$ .



**Figure 6.** Products from microwave heating (17 min at 180°C) with standard triethylamine concentration. Figure 6A is from a microemulsion with composition A' on Figure 1. Figure 6B is the control experiment without the microemulsion. The scale bars are A-2  $\mu\text{m}$  and B-20  $\mu\text{m}$ .



**Figure 7.** AlPO<sub>4</sub>-5 fibers crystallized by microwave heating (17 min at 180°C) with double the standard amount of triethylamine from a microemulsion with the weight ratio of CPC to butanol is 3 to 1. The scale bar is 2 microns.



A




B



C




D




E

FIG. 7






**FIG. 8**



**FIG. 9**



**FIG. 10**




FIG. 11




FIG. 12